

A NATIONAL NORM FOR THE COUNT VARIATION OF YARN PRODUCED IN

SRI LANKA

by



CYNTHIA WANNIARACHCHI

www.lib.mrt.ac.lk

LIBRARY
UNIVERSITY OF MORATUWA, SRI LANKA
MORATUWA



A dissertation submitted to the UNIVERSITY OF MORATUWA in
part fulfilment of the POST GRADUATE DIPLOMA IN
TEXTILE TECHNOLOGY

677.074 (548.7)

Department of Textile Technology,
University of Moratuwa,
Moratuwa,
SRI LANKA.

Department of Textile Technology

University of Moratuwa



78497

1983

78497

78497

CONTENTS

Acknowledgements

Abstract

CHAPTER 1 - Introduction	page

1.1 Yarn count variation	1
1.2 Literature survey	3
CHAPTER 2 - Effects of yarn count variation on Textiles	11

CHAPTER 3 - Experimental	

3.1 Samples used for measurements	14
3.2 Sampling	14
3.3 Method	14
3.4 Results of yarn count measurements	15
3.5 Statistical analysis of results	15
CHAPTER 4 - Conclusion	16

APPENDICES	

APPENDIX A	19
APPENDIX B	24
LITERATURE CITED	27

ACKNOWLEDGEMENTS

Sincere thanks are extended to the assistance of Mr. Jack Woolfenden of the Textile Training and Services Centre, and Mr. Rohana Kuruppu of the University of Moratuwa in developing the particular approach to this project.

In addition, I express appreciation for the assistance received from the other members of the staff of the University of Moratuwa and the Textile Training and Services Centre.

Thanks are also expressed to the respective managers of mills for permitting me to collect samples and to use them in this project.

I also wish to record my appreciation of the comments, suggestions, and other help received from my good friends.

Last but not least I thank the Sri Lanka Standards Institution for allowing me to undertake this valuable course.

ABSTRACT

This project is aimed at establishing a National norm for the count variation of yarns produced in Sri Lankan mills. Owing to, (a) its simplicity (b) its ease of understanding by most mill personnel (c) the possibility of making some international comparisons on National norms and (d) the ease by which data can be collected, the Percentage Mean Range (PMR) method of measuring count variation has been used in this project.



University of Moratuwa, Sri Lanka.
Electronic Theses & Dissertations
www.lib.mrt.ac.lk

The results of this project have indicated that the average count variation of Sri Lankan spun carded cotton yarns are very much greater than (worse than) the world average value given under USTER STATISTICS 1982.